Author Index

Abe, K., Kawagoe, J., Lee, T.-H., Aoki, M. and Kogure, K.
Disturbance of a mitochondrial DNA expression in gerbil hippocampus after transient forebrain ischemia (19) 69
Aceves, J., see Arias-Montaño, J.-A. (19) 233
Agarwal, N., see Nir, I. (19) 47
Akagawa, K., see Inoue, A. (19) 121
Alkon, D.L., see Craig, A.M. (19) 269
Aoki, M., see Abe, K. (19) 69
Arias-Montaño, J.-A., Aceves, J. and Young, J.M.
Carbachol-induced phosphoinositide metabolism in slices of rat substantia nigra pars reticulata (19) 233
Asanuma, M., see Chou, H.-h. (19) 211
Azmitia, E.C., see Liao, B. (19) 328

Baldessari, S., see Bendotti, C. (19) 257
Barizo, M.J., see Sanna, P.P. (19) 241
Baudry, M., see Wei, W. (19) 339
Bendotti, C., Baldessari, S., Ehret, M., Tarizzo, G. and Samanin, R.
Effect of d-fenfluramine and 5,7-dihydroxytryptamine on the levels of tryptophan hydroxylase and its mRNA in rat brain (19) 257
Bengzon, J., see Kokaia, Z. (19) 277
Bloom, F.E., see Sanna, P.P. (19) 241
Bogic, L., see Hammer Jr., R.P. (19) 129
Bourbonnière, M. and Nalbantoglu, J.
Expression of amyloid precursor protein in a neuronal cell line: functional activity of proximal regulatory elements (19) 246
Bredesen, D.E., see Zhong, L.-t. (19) 353
Broek, D., see Wei, W. (19) 339
Brooks, P.J., Funabashi, T., Kleopoulos, S.P., Mobbs, C.V. and Pfaff, D.W.
Cell-specific expression of preproenkephalin intronic heteronuclear RNA in the rat forebrain (19) 22
Burgess, L.H. and Handa, R.J.
Hormonal regulation of androgen receptor mRNA in the brain and anterior pituitary gland of the male rat (19) 31
Byrne, J.H., see Noel, F. (19) 203

Chou, H.-h., Ogawa, N., Asanuma, M., Hirata, H., Kondo, Y. and Mori, A.
Rapid response of striatal muscarinic M_1-receptor mRNA to muscarinic cholinergic agents in rat brain (19) 211
Chritin, M., Feuerstein, C. and Savasta, M.
Time-course of changes in striatal levels of DA uptake sites, DA D_2 receptor and preproenkephalin mRNAs after nigrostriatal dopaminergic denervation in the rat (19) 318
Chua Jr., S.C., see Jhanwar-Uniyal, M. (19) 195
Cook, R., see Noel, F. (19) 203
Craig, A.M., Olds, J.L., Schreurs, B.G., Scharenberg, A.M. and Alkon, D.L.
Quantitative distribution of protein kinase C α, β, γ, and ϵ mRNAS in the hippocampus of control and nictitating membrane conditioned rabbits (19) 269

De Blas, A., see Luntz-Leybman, V. (19) 9
De Groen, P.C., see Nielander, H.B. (19) 293
Devi, L., see Grigoriants, O. (19) 161
Donnelly-Roberts, D.L. and Lentz, T.L.
Sodium dodecyl sulfate- and carbamylcholine-induced changes in circular dichroism spectra of acetylcholine receptor synthetic peptides (19) 55
Dreyer, D., see Pennypacker, K.R. (19) 349
Duman, R.S., see Vaccarino, F.M. (19) 76

Eggen, B.J.L., see Nielander, H.B. (19) 293
Ehret, M., see Bendotti, C. (19) 257
Elizarov, S.M. and Preobrazhensky, A.A.
Phosphorylation of two human neurochordins by mammalian casein kinase 1 (19) 310
Elliott, K., see Franklin, S.O. (19) 93
Eskin, A., see Noel, F. (19) 203

Fei, J., see Lam, D.M.-K. (19) 227
Feng, P., see Satoh, T. (19) 175
Feng, X., see Katagiri, T. (19) 1
Fernandez, J., see Schuligoi, R. (19) 333
Fernando, L., see Luntz-Leybman, V. (19) 9
Feuerstein, C., see Chritin, M. (19) 318
Fischer-Colbrie, R., see Mahata, S.K. (19) 83
Folsom, D.P., see Sanna, P.P. (19) 241
Franklin, S.O., Elliott, K., Zhu, Y.-S., Wahlestedt, C. and Inturrisi, C.E.
Quantitation of NMDA receptor (NMDAR1) mRNA levels in the adult and developing rat CNS (19) 93
Fricker, L.D., see Grigoriants, O. (19) 161
Frostholm, A., see Luntz-Leybman, V. (19) 9
Fujita, M., Shimada, S., Nishimura, T., Uhl, G.R. and Tohyama, M.
Ontogeny of dopamine transporter mRNA expression in the rat brain (19) 222
Funabashi, T., see Brooks, P.J. (19) 22
Furukawa, T., see Shibata, K. (19) 115
Furuyama, T., Inagaki, S. and Takagi, H.
Distribution of type II adenylyl cyclase mRNA in the rat brain (19) 165

Gidö, G., see Kokaia, Z. (19) 277
Gispen, W.H., see Nielander, H.B. (19) 293
Gray, C.W. and Patel, A.J.
Regulation of β-amyloid precursor protein isoform mRNAs by transforming growth factor-β1 and interleukin-1β in astrocytes (19) 251
Grigoriants, O., Devi, L. and Fricker, L.D.
Dopamine antagonist haloperidol increases carboxypeptidase E mRNA in rat neurointermediate pituitary but not in various other rat tissues (19) 161
Guo, L.-H., see Lam, D.M.-K. (19) 227

Hammer Jr., R.P., Bogic, L. and Handa, R.J.
Estrogenic regulation of proenkephalin mRNA expression in the ventromedial hypothalamus of the adult male rat (19) 129
Handa, R.J., see Burgess, L.H. (19) 31
Handa, R.J., see Hammer Jr., R.P. (19) 129
Hartigan, D.J., see Vaccarino, F.M. (19) 76
Hayward, M.D., see Vaccarino, F.M. (19) 76
Heavens, R.P., see Schuligoi, R. (19) 333
Hirata, H., see Chou, H.-h. (19) 211
Hirsch, M.D., see Sanna, P.P. (19) 241
Hockfield, S., see Martin, K.A. (19) 303
Hong, J.S., see Pennypacker, K.R. (19) 349
Horn, D.A. and Latchman, D.S.
The tissue specific SmN protein does not influence the alternative splicing of endogenous N-Cam and C-SRC RNAs in transfected 3T3 cells (19) 181
Huang, F., see Lam, D.M.-K. (19) 227

Iadarola, M.J., see Jian, M. (19) 156
Ichikawa, T., see Katagiri, T. (19) 1
Ihara, Y., see Kobayashi, H. (19) 188
Inagaki, S., see Furuyama, T. (19) 165
Inoue, A. and Akagawa, K.
Neuron specific expression of a membrane protein, HPC-1: tissue distribution, and cellular and subcellular localization of immunoreactivity and mRNA (19) 121
Inturrisi, C.E., see Franklin, S.O. (19) 93

Jhanwar-Uniyal, M. and Chua Jr., S.C.
Critical effects of aging and nutritional state on hypothalamic neuropeptide Y and galanin gene expression in lean and genetically obese Zucker rats (19) 195
Jian, M., Staines, W.A., Iadarola, M.J. and Robertson, G.S.
Destruction of the nigrostriatal pathway increases Fos-like immunoreactivity predominantly in striatopallidal neurons (19) 156
Johnson, G., Refolo, L.M., Merril, C.R. and Wallace, W.
Altered expression and phosphorylation of amyloid precursor protein in heat shocked neuronal PC12 cells (19) 140
Johnson, G., see Wallace, W. (19) 149

Kalivas, P.W., see See, R.E. (19) 219
Kane, D.J., see Zhong, L.-t. (19) 353
Katagiri, T., Feng, X., Ichikawa, T., Usui, H., Takahashi, Y. and Kumanishi, T.
Neuron-specific enolase (NSE) and non-neuronal enolase (NNE) mRNAs are co-

expressed in neurons of the rat cerebellum: in situ hybridization histochemistry (19) 1
Kawagoe, J., see Abe, K. (19) 69
Kawasaki, M., Yoshihara, Y., Yamaji, M. and Watanabe, Y.
Expression of prostaglandin endoperoxide synthase in rat brain (19) 39
Kim, U.J., see Satoh, T. (19) 175
King, S.C., see Lam, D.M.-K. (19) 227
Kiyama, H., Sato, K. and Tohyama, M.
Characteristic localization of non-NMDA type glutamate receptor subunits in the rat pituitary gland (19) 262
Kiyama, H., Sato, K., Kuba, T. and Tohyama, M.
Sympathetic and parasympathetic ganglia express non-NMDA type glutamate receptors: distinct receptor subunit composition in the principle and SIF cells (19) 345
Kiyama, H., see Saika, T. (19) 237
Kleopoulos, S.P., see Brooks, P.J. (19) 22
Kobayashi, H., Uchida, Y., Ihara, Y., Nakajima, K., Kohsaka, S., Miyatake, T. and Tsuji, S.
Molecular cloning of rat growth inhibitory factor cDNA and the expression in the central nervous system (19) 188
Kogure, K., see Abe, K. (19) 69
Kohsaka, S., see Kobayashi, H. (19) 188
Kojima, N., Shirao, T. and Obata, K.
Molecular cloning of a developmentally regulated brain protein, chicken drebrin A and its expression by alternative splicing of the drebrin gene (19) 101
Kokaia, M., see Kokaia, Z. (19) 277
Kokaia, Z., Gidö, G., Ringstedt, T., Bengzon, J., Kokaia, M., Siesjö, B.K., Persson, H. and Lindvall, O.
Rapid increase of BDNF mRNA levels in cortical neurons following spreading depression: regulation by glutamatergic mechanisms independent of seizure activity (19) 277
Komatsu, C., see Shibata, K. (19) 115
Kondo, H., see Sakagami, H. (19) 215
Kondo, Y., see Chou, H.-h. (19) 211
Kreek, M.J., see Spangler, R. (19) 323
Krukoff, T.L. and Zheng, Y.
Effects of renal denervation and reinnervation on ganglionic gene expression of neurotransmitter proteins and c-*fos* in rat (19) 287
Kuba, T., see Kiyama, H. (19) 345
Kubo, T., see Saika, T. (19) 237
Kumanishi, T., see Katagiri, T. (19) 1

Lam, D.M.-K., Fei, J., Zhang, X.-Y., Tam, A.C.W., Zhu, L.-H., Huang, F., King, S.C. and Guo, L.-H.
Molecular cloning and structure of the human (GABATHG) GABA transporter gene (19) 227
Latchman, D.S., see Horn, D.A. (19) 181
Le, H.N., see Vaccarino, F.M. (19) 76
Lee, T.-H., see Abe, K. (19) 69
Lentz, T.L., see Donnelly-Roberts, D.L. (19) 55
Liao, B., Miesak, B. and Azmitia, E.C.
Loss of 5-HT_{1A} receptor mRNA in the dentate gyrus of the long-term adrenalectomized rats and rapid reversal by dexamethasone (19) 328
Lindvall, O., see Kokaia, Z. (19) 277
Luntz-Leybman, V., Frostholm, A., Fernando, L., De Blas, A. and Rotter, A.
$GABA_A$/benzodiazepine receptor γ_2 subunit gene expression in developing normal and mutant mouse cerebellum (19) 9

Maciejewski-Lenoir, D., see Sanna, P.P. (19) 241
Maggi, A., see Pollio, G. (19) 135
Mahata, M., see Mahata, S.K. (19) 83
Mahata, S.K., Mahata, M., Fischer-Colbrie, R. and Winkler, H.
Reserpine causes differential changes in the mRNA levels of chromogranin B, secretogranin II, carboxypeptidase H, alpha-amidating monooxygenase, the vesicular amine transporter and of synaptin/synaptophysin in rat brain (19) 83
Martin, K.A. and Hockfield, S.
Expression of the *mas* proto-oncogene in the rat hippocampal formation is regulated by neuronal activity (19) 303
Matsunaga, T., see Saika, T. (19) 237
McMillian, M.K., see Pennypacker, K.R. (19) 349
Melia, K.R., see Sanna, P.P. (19) 241
Merril, C.R., see Johnson, G. (19) 140
Merril, C.R., see Wallace, W. (19) 149
Miesak, B., see Liao, B. (19) 328
Misumi, Y., see Shibata, K. (19) 115
Miyatake, T., see Kobayashi, H. (19) 188
Miyatake, T., see Saito, F. (19) 171
Mobbs, C.V., see Brooks, P.J. (19) 22
Morgan, J.I., see Sangameswaran, L. (19) 62
Mori, A., see Chou, H.-h. (19) 211

Nakajima, K., see Kobayashi, H. (19) 188
Nalbantoglu, J., see Bourbonnière, M. (19) 246
Nestler, E.J., see Vaccarino, F.M. (19) 76
Nicolin, A., see Pollio, G. (19) 135
Nielander, H.B., De Groen, P.C., Eggen, B.J.L., Schrama, L.H., Gispen, W.H. and Schotman, P.
Structure of the human gene for the neural phosphoprotein B-50 (GAP-43) (19) 293
Nir, I. and Agarwal, N.
Diurnal expression of c-*fos* in the mouse retina (19) 47
Nishimura, T., see Fujita, M. (19) 222
Noel, F., Nuñez-Regueiro, M., Cook, R., Byrne, J.H. and Eskin, A.
Long-term changes in synthesis of intermediate filament protein, actin and other proteins in pleural sensory neurons of *Aplysia* produced by an in vitro analogue of sensitization training (19) 203
Nuñez-Regueiro, M., see Noel, F. (19) 203

Obata, K., see Kojima, N. (19) 101
Ogawa, N., see Chou, H.-h. (19) 211
Olds, J.L., see Craig, A.M. (19) 269

Patel, A.J., see Gray, C.W. (19) 251
Pennypacker, K.R., Dreyer, D., Hong, J.S. and McMillian, M.K.
Elevated basal AP-1 DNA binding activity in developing rat brain (19) 349
Persson, H., see Kokaia, Z. (19) 277
Pfaff, D.W., see Brooks, P.J. (19) 22
Pollio, G., Xue, P., Zanisi, M., Nicolin, A. and Maggi, A.
Antisense oligonucleotide blocks progesterone-induced lordosis behavior in ovariectomized rats (19) 135
Preobrazhensky, A.A., see Elizarov, S.M. (19) 310

Refolo, L.M., see Johnson, G. (19) 140
Refolo, L.M., see Wallace, W. (19) 149
Ringstedt, T., see Kokaia, Z. (19) 277
Robertson, G.S., see Jian, M. (19) 156
Rotter, A., see Luntz-Leybman, V. (19) 9

Saika, T., Takeda, N., Kiyama, H., Kubo, T., Tohyama, M. and Matsunaga, T.
Changes in preproenkephalin mRNA after unilateral and bilateral labyrinthectomy in the rat medial vestibular nucleus (19) 237
Saito, F., Yanagisawa, K. and Miyatake, T.
Soluble derivatives of β/A4 amyloid protein precursor in human cerebrospinal fluid are both *N*- and *O*-glycosylated (19) 171
Sakagami, H. and Kondo, H.
Cloning and sequencing of a gene encoding the β polypeptide of Ca^{2+}/calmodulin-dependent protein kinase IV and its expression confined to the mature cerebellar granule cells (19) 215
Samanin, R., see Bendotti, C. (19) 257
Sangameswaran, L. and Morgan, J.I.
Structure and regulation of the gene encoding the neuron-specific protein PEP-19 (19) 62
Sanna, P.P., Folsom, D.P., Barizo, M.J., Hirsch, M.D., Melia, K.R., Maciejewski-Lenoir, D. and Bloom, F.E.
Chronic ethanol intake decreases vasopressin mRNA content in the rat hypothalamus: a PCR study (19) 241
Sato, K., see Kiyama, H. (19) 262
Sato, K., see Kiyama, H. (19) 345
Satoh, T., Feng, P., Kim, U.J. and Wilber, J.F.
Identification of thyrotropin-releasing hormone receptor messenger RNA in the rat central nervous system and eye (19) 175
Savasta, M., see Chritin, M. (19) 318
Scharenberg, A.M., see Craig, A.M. (19) 269
Schotman, P., see Nielander, H.B. (19) 293
Schrama, L.H., see Nielander, H.B. (19) 293
Schreiber, S.S., see Wei, W. (19) 339
Schreurs, B.G., see Craig, A.M. (19) 269
Schuligoi, R., Fernandez, J., Heavens, R.P. and Sirinathsinghji, D.J.S.

Decreased tyrosine hydroxylase mRNA but not cholecystokinin mRNA in the pars compacta of the substantia nigra and ventral tegmental area of aged rats (19) 333
Sealfon, S.C., see Spyraki, C. (19) 313
See, R.E., Striplin, C. and Kalivas, P.W.
Chronic haloperidol does not alter G protein α-subunit levels in rats (19) 219
Shibata, K., Komatsu, C., Misumi, Y. and Furukawa, T.
Developmental differences of angiotensinogen mRNA in the preoptic area between spontaneously hypertensive and age-matched Wistar-Kyoto rats (19) 115
Shimada, S., see Fujita, M. (19) 222
Shirao, T., see Kojima, N. (19) 101
Siesjö, B.K., see Kokaia, Z. (19) 277
Sirinathsinghji, D.J.S., see Schuligoi, R. (19) 333
Spangler, R., Unterwald, E.M. and Kreek, M.J.
'Binge' cocaine administration induces a sustained increase of prodynorphin mRNA in rat caudate-putamen (19) 323
Spyraki, C. and Sealfon, S.C.
Regulation of dopamine D_2 receptor mRNA expression in the olfactory tubercle by cocaine (19) 313
Staines, W.A., see Jian, M. (19) 156
Striplin, C., see See, R.E. (19) 219
Sugar, J., see Wallace, W. (19) 149

Takagi, H., see Furuyama, T. (19) 165
Takahashi, Y., see Katagiri, T. (19) 1
Takeda, N., see Saika, T. (19) 237
Tam, A.C.W., see Lam, D.M.-K. (19) 227
Tarizzo, G., see Bendotti, C. (19) 257
Tocco, G., see Wei, W. (19) 339
Tohyama, M., see Fujita, M. (19) 222
Tohyama, M., see Kiyama, H. (19) 262
Tohyama, M., see Kiyama, H. (19) 345
Tohyama, M., see Saika, T. (19) 237
Tsuji, S., see Kobayashi, H. (19) 188

Uchida, Y., see Kobayashi, H. (19) 188
Uhl, G.R., see Fujita, M. (19) 222
Unterwald, E.M., see Spangler, R. (19) 323
Usui, H., see Katagiri, T. (19) 1

Vaccarino, F.M., Hayward, M.D., Le, H.N., Hartigan, D.J., Duman, R.S. and Nestler, E.J.
Induction of immediate early genes by cyclic AMP in primary cultures of neurons from rat cerebral cortex (19) 76

Wahlestedt, C., see Franklin, S.O. (19) 93
Wallace, W., Johnson, G., Sugar, J., Merril, C.R. and Refolo, L.M.
Reversible phosphorylation of tau to form A68 in heat-shocked neuronal PC12 cells (19) 149
Wallace, W., see Johnson, G. (19) 140
Watanabe, Y., see Kawasaki, M. (19) 39
Wei, W., Schreiber, S.S., Baudry, M., Tocco, G. and Broek, D.
Localization of the cellular expression pattern of *cdc25*^{NEF} and *ras* in the juvenile rat brain (19) 339
Wilber, J.F., see Satoh, T. (19) 175
Winkler, H., see Mahata, S.K. (19) 83

Xue, P., see Pollio, G. (19) 135

Yamaji, M., see Kawasaki, M. (19) 39
Yanagisawa, K., see Saito, F. (19) 171
Yoshihara, Y., see Kawasaki, M. (19) 39
Young, J.M., see Arias-Montaño, J.-A. (19) 233

Zanisi, M., see Pollio, G. (19) 135
Zhang, X.-Y., see Lam, D.M.-K. (19) 227
Zheng, Y., see Krukoff, T.L. (19) 287
Zhong, L.-t., Kane, D.J. and Bredesen, D.E.
BCL-2 blocks glutamate toxicity in neural cell lines (19) 353
Zhu, L.-H., see Lam, D.M.-K. (19) 227
Zhu, Y.-S., see Franklin, S.O. (19) 93